

A Charged Fusion Product Diagnostic for a Spherical Tokamak


Proposal Defense
Ramona V Perez

Outline

1. Background
2. Experimental Design
 1. Mechanical Design
 2. Simulations
 3. Electronic & Data Acquisition Design
3. Data Collection
4. Data Analysis
5. Timeline



Confined Plasma

- Poloidal & toroidal direction
- Temperature 10^8 K
- Magnetic field .5T




[Image 1]

Confined Plasma



[Image 3]


Spherical Tokamak




Aspect Ratio: R/a

[Image 4]

Tokamak Field Coils



Heating Methods




1. Ohmic heating
(induced current)


2. **Neutral-Beam
Injection (NBI)**

And Radio Frequency (RF)
(oscillating
electromagnetic waves)


Products and Reactants




Proton
(Hydrogen
nucleus) **P**




Deuterium nucleus
(Hydrogen isotope) **D**



Tritium nucleus
(Hydrogen isotope) **T**



Helium3 nucleus
(Helium isotope) **He³**



Helium4 nucleus (Helium
isotope) **α**

Fusion Reactions

- Primary reactions



- Secondary reactions



Detect these



DD Reaction

- Energy investment to facilitate nuclear reactions





Figure 3.11 Velocity averaged cross section (i.e., $\langle \sigma v \rangle = R_{ij}/n_i n_j$) for the D-T, D-He³, and D-D fusion reactions as a function of temperature.

Research Objective


- Determine emission profile
 - Where are the $d(d,p)t$ reactions taking place in the plasma?
 - At what rate are these $d(d,p)t$ reactions taking place in the plasma?
- MHD instabilities
- Provides foundation for future work with spherical tokamaks

NSTX and MAST






Tokamak

[Image 5]




[Image 6]



Module Cross Section




Total Exploded View




Total Assembled View





MAST mid-plane
cross-sectional
top view



Reciprocating probe


Attach
diagnostic

[Image 11]

2.1 Experimental Design. Mechanical Design

MAST RP Access Cube

- Clearance Diameter: 148mm
- CFPD Diameter: 111mm
- CFPD Length: 201.7mm
= 185mm + 1.7cm





2.2 Simulations

- Orbit Code
- Recreate particle trajectory, or orbit, backwards in time

$$\epsilon_{sim} = \frac{\sum_{n=1}^{N_{det}} S_n}{\sum_{n=1}^{N_{total}} S_n}$$

$S_n = \text{emissivity for event } n$



Flux Surfaces

- Constant pressure
- Constant temperature

$$\epsilon_{sim} = \frac{\sum_{n=1}^{N_{det}} S_n}{\sum_{n=1}^{N_{total}} S_n}$$

$S_n = \text{emissivity for event } n$








Figure 11.8 Contours of constant pressure in a well-confined toroidal equilibrium.



[Image 15]


Figure 11.9 Two flux surfaces, 1 and 2, at two different toroidal locations showing that the current flows between and not across them.

Particle Orbits





Other MAST Diagnostics

- All diagnostics on the tokamak study the same plasma





2.3 Electronics & DAQ Design







MAST RP Access Cube

- MAST will connect non terminated cable ends from detectors to RP connector



MAST RP Linkbox


- 4 preamplifiers stored inside linkbox
- MAST will connect non terminated cable ends from preamplifiers to RP
- Bias supply and power supply cables to preamp will run into linkbox



[Image 18]

Hardware Storage

- 10m from MAST RP
 - 4 amplifiers
 - 1 NIM BIN
 - 1 rack mount computer
 - 1 PCI extension box
 - 4 power supplies for detector bias Voltage



Data Files (HDF)

- Data file sizes
 - 23MB per channel per shot
 - 92MB per shot for all channels
 - 46GB for 10 days (2 weeks) data collection
 - 92GB for 4 weeks data collection
- Data Storage
 - 150GB onsite rackmount computer
 - 800GB FIU host computer

The screenshot displays a software interface for viewing an HDF5 file. On the left, a tree view shows the file structure for '500ms_25V.hws'. The tree includes folders for 'cfg_scope0', 'wfm_group0', 'axes', 'id', 'traces', and 'vectors'. Under 'traces', there is a folder 'trace0' which contains sub-folders for 'configs', 'x-axis', and 'y-axis'. The 'y-axis' folder contains a 'data_vector' folder, which in turn contains a 'data' dataset and a 'scale_coef' dataset.

On the right, a 'TableView' window shows a table of data. The table has two columns: an index from 0 to 41 and a corresponding numerical value. The values range from -128 to -160, with some positive values like 32 and 80.

| Index | Value |
|-------|-------|
| 0 | -128 |
| 1 | -112 |
| 2 | -80 |
| 3 | -96 |
| 4 | -80 |
| 5 | -64 |
| 6 | -80 |
| 7 | -112 |
| 8 | -144 |
| 9 | -144 |
| 10 | -112 |
| 11 | -96 |
| 12 | -64 |
| 13 | -16 |
| 14 | -48 |
| 15 | -112 |
| 16 | -80 |
| 17 | -32 |
| 18 | -48 |
| 19 | -48 |
| 20 | 32 |
| 21 | 80 |
| 22 | 0 |
| 23 | -16 |
| 24 | 16 |
| 25 | -16 |
| 26 | -32 |
| 27 | 16 |
| 28 | -32 |
| 29 | -128 |
| 30 | -64 |
| 31 | 16 |
| 32 | -112 |
| 33 | -176 |
| 34 | -80 |
| 35 | -32 |
| 36 | -112 |
| 37 | -128 |
| 38 | -32 |
| 39 | -64 |
| 40 | -176 |
| 41 | -160 |


At the bottom of the interface, a metadata panel for the 'data' dataset is visible. It shows the dataset name 'data (315032)', its data type '16-bit integer', a size of '30000000', and 'Number of attributes = 0'. There are tabs for 'Log Info' and 'Metadata'.

3. Data Collection: *Round 2*



- Initial noise diagnostics and Electrical Design Review
 - January/March 2013
 - The Mega Amp Spherical Tokamak (MAST) at in the Culham Centre for Fusion Energy (CCFE) in the United Kingdom
- Diagnostic installation and subsequent data collection
 - May/ June 2013
 - The Mega Amp Spherical Tokamak (MAST) at in the Culham Centre for Fusion Energy (CCFE) in the United Kingdom

4. Data Analysis


- Example of pulse signals, 5.5 MeV alpha particles




Find Peaks for Protons





Bin Peaks




Histogram 5 for Peak Values



Count Particles Detected



Emissivity



- Ratio of particles detected (counted) and MAST global yield rate proportional to below expression
- Simple models of emissivity will be fitted to data

$$\epsilon = \frac{\int A(\theta) d\theta \int_{orbit} S(\vec{r}) dl}{2\pi \int_V S(\vec{r}) dV}$$

$S(\vec{r}) = \text{emissivity at position } \vec{r}$

32 $A(\theta) = \text{effective detector opening for entry angle } \theta$

5. Tentative Timeline

| TERM | GOALS |
|-------------|--|
| SUMMER 2012 | Diagnostic design/ assembly/ testing |
| FALL 2012 | Continue summer goal, apply for 2014 fellowship |
| SPRING 2013 | Initial testing & Electrical Design Review at MAST |
| SUMMER 2013 | Diagnostic installation and data collection |
| FALL 2013 | Data analysis, dissertation, apply for 2015 fellowship |
| SPRING 2014 | Data analysis, dissertation |
| SUMMER 2014 | Write paper for publication, dissertation |
| FALL 2014 | Dissertation defense, submit paper for publication |

References- Images

- Unreferenced images are created by author
- 1. An image depicting the poloidal (red, called theta) direction and the toroidal (blue, called phi) directions. 13 September 2006. Made in POV-Ray by Dave Burke. PNG File.
- 2. European Fusion Development Agreement (EFDA). Magnetic fields in a tokamak. 2012. Garching, Germany. EFDA: Fusion. 08/01/2012. <<http://www.efda.org/fusion/focus-on/plasma-heating-current-drive/ohmic-heating/>>
- 3. Jeffrey Freidberg. Cover. Plasma Physics and Fusion Energy. Cambridge University Press, 2008.
- 4. Spherical Tokamaks Image. 2009. Abingdon, Oxfordshire, UK. Culham Centre for Fusion Energy: Research. 08/01/2012. <<http://www.ccf.ac.uk/st.aspx>>
- 5. Princeton Plasma Physics Laboratory. National Spherical Torus Experiment. Princeton, New Jersey. Alternative Energy Action Now. 09/01/2012. <<http://www.alternative-energy-action-now.com/spherical-tokamak.html>>
- 6. Spherical Tokamaks Image. 2009. Abingdon, Oxfordshire, UK. Culham Centre for Fusion Energy: Research. 08/01/2012. <<http://www.ccf.ac.uk/st.aspx>>
- 10. Tokamak Fields. Encyclopedia of Earth. 09/01/2012. [imagehttp://www.eoearth.org/article/Nuclear_fusion_power](http://www.eoearth.org/article/Nuclear_fusion_power).
- 6. European Fusion Development Agreement (EFDA). Heating of JET Plasmas. 2012. Garching, Germany. EFDA: Fusion. 08/01/2012. <http://www.efda.org/fusion/focus-on/plasma-heating-current-drive/ohmic-heating/>
- 7. Fig. 1. Fundamental quantum mechanical phenomena. 2012. Victoria Stafforda Psychic Investigation. 09/20/2012. <<http://victoriastaffordapsychicinvestigation.wordpress.com/2012/07/01/line-19-a2a-semiconductor-heterostructures-schrodinger-quantum-confinement-5g-wow-seti/fig-1-fundamental-quantum-mechanical-phenomena-a-electron-reflection-and-interference-b-tunneling-effect-c-e-quantum-confinement/>>

References- Images

10. Figure 3.11. Page 51, 2007. Plasma Physics and Fusion Energy. Jeffrey Freidberg. Cambridge University Press, 2007.
11. Side View of Assembled MAST Reciprocating Probe. CAD image created by the Culham Centre for Fusion Energy's MAST Drawing Office. 2012. PDF File.
12. MAST Reciprocating Probe Access Cube. CAD image created by the Culham Centre for Fusion Energy's MAST Drawing Office. 2012. PDF File.
13. Side View of MAST Reciprocating Probe. CAD image created by the Culham Centre for Fusion Energy's MAST Drawing Office. 2012. PDF File.
14. Figure 11.8 and Figure 11.9. Page 262, 2007. Plasma Physics and Fusion Energy. Jeffrey Freidberg. Cambridge University Press, 2007.
15. Cecconello *et al.* FIG 2. MAST Neutron camera schematic. Rev Sci Instrum. **81**, 10D315 (2010).
17. Scott Y. Allan. MAST Reciprocating Probe. Culham Centre for Fusion Energy. 2012. JPG File.
18. Linkbox MAST Reciprocating Probe. CAD image created by the Culham Centre for Fusion Energy's MAST Drawing Office. 2012. PDF File.
19. Please ask for user manual, CANBERRA 2111 Timing Filter Amplifier.
20. Please ask for user manual, ADNACO S2 Fiber Optic PCI Bus Extender.
21. Please ask for user manual, SuperMicro 50161 MTF 1U Rackmount Server.
22. Please ask for user manual, CANBERRA 3002D 0-3 kV H.V. Power Supply.
23. Please ask for user manual, Tennelec TB-3 NIM BIN with TC-911 Power Supply System.


References

1. J. A. Bittencourt, *FUNDAMENTALS OF PLASMA PHYSICS*. Springer Science + Business Media, LLC, 3rd Edition, 2004.
2. W. U. Boeglin, R. Valenzuela Perez, D. S. Darrow, Rev. Sci. Instrum. **81** (2010) 10D301
3. Hans-Stephan Bosch, Rev. Sci. Instrum. **61**, 1699 (1990)
4. L. F. Delgado-Aparicio, et. al., J. of Appl. Phys. **102** (2007) 073304
5. Jeffrey Freidberg, *Plasma Physics and Fusion Energy*. Cambridge University Press, 2007.
6. Daniel H. Lo, Réjean L. Boivin, and Richard D. Petrasso Rev. Sci. Instrum. **66**, 345 (1995)
7. J. D. Strachan, Rev. Sci. Instrum. **57**, 1771 (1986)
8. S. J. Zweben, Rev. Sci. Instrum. **57**, 1774 (1996)
9. S. J., Zweben, et al., Nucl. Fusion **35**, 893 (1995)


Thank you for your time!

Questions

Alternate Washer to Change Collimator Size




New collimator size




Particle Motion

- How do particles move in these magnetic fields?

$$Radius_{gyro} = \frac{mv_{\perp B}}{|q|B}$$






Tokamak

- Facilitate nuclear reactions

$$Potential_{Coulomb} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r}$$



Gyro radius

$$Radius_{gyro} = \frac{mv_{\perp B}}{|q|B}$$

$$m \frac{dv_{\perp B}}{dt} = qv_{\perp B} \times B$$

$$m \frac{dv_{\perp B}}{dt} = \frac{mv_{\perp B}^2}{Radius_{gyro}}$$

Collimator





Fig. 1. Solid angle subtended by a detector of area D without (right) and with a cylindrical (middle) and a conical (left) collimator.

[Image #]


Transformer



Copyright UOregon

[Image #]

Electronics Schema



Module Exploded View with Bases

